

AQUACEL™ Extra™ e AQUACEL™ Ag Extra™ gestiscono alti livelli di essudato



Scegli la tua soluzione

Misure (cm)	N. pezzi	Codice Prodotto	Codice ISO
AQUACEL™ Extra™			
5 x 5 cm	10	420671	
10 x 10 cm	10	420672	09.21.12.003
15 x 15 cm	5	420673	
AQUACEL™ Ag Extra™			
5 x 5 cm	10	420675	
10 x 10 cm	10	420676	09.21.12.003
15 x 15 cm	5	420678	
20 x 30 cm	5	420679	
AQUACEL™ NASTRO con fibra rinforzante			
2 x 45 cm	5	403770	
1 x 45 cm	5	420127	
AQUACEL™ Ag NASTRO con fibra rinforzante			
2 x 45 cm	5	403771	
1 x 45 cm	5	420128	

AQUACEL™ Extra™	CND M040404 - M040413	N. di repertorio: 445834 - 445895
AQUACEL™ Ag Extra™	CND M040404 - M040413 - M04040802	N. di repertorio: 645328 - 645327 - 645252
AQUACEL™ NASTRO con fibra rinforzante	CND M040404 - M040413	N. di repertorio: 6504-176807
AQUACEL™ Ag NASTRO con fibra rinforzante	CND M040404 - M040413 - M04040802	N. di repertorio: 70728

Bibliografia: 1. World Union of Wound Healing Societies (WUWHS). *Principles of best practice: wound exudate and the role of dressings. A consensus document*. London: MEP Ltd. 2007. 2. Newman GR, Walker M, Hobot JA, Bowler PG. 2006. Visualisation of bacterial sequestration and bacterial activity within hydrating Hydrofiber® wound dressings. *Biomaterials* 27(10/1): 1129-1139. 3. Walker M, Hobot JA, Newman GR, Bowler PG. 2003. Scanning electron microscopic examination of bacterial immobilization in a carboxymethyl cellulose (AQUACEL™) and alginate Dressing. *Biomaterials* 24:883-890. 4. Bowler PG, Jones SA, Davies RJ, Coyne E. 1999. Infection control properties of some wound dressings. *J Wound Care* 8(10):499-502. 5. Walker M, Bowler PG, Cochrane CA. 2007. In vitro studies to show sequestration of matrix metalloproteinases by silver-containing wound care products. *Ostomy/Wound Management* 53(8): 19-25. 6. Walker M and Parsons D. 2010. Hydrofiber™ Technology: its role in exudate management. *Wounds UK* 6: 31-38. 7. Parsons D, Bowler PG, Myles V, Jones SA. 2005. Silver antimicrobial dressings in wound management: A comparison of antibacterial, physical and chemical characteristics. *WOUNDS*, 17: 222-232. 8. Jones SA, Bowler PG, Walker M. 2005. Antimicrobial activity of silver-containing dressings is influenced by dressing conformability with a wound surface. *WOUNDS*, 17: 263-270. 9. Bishop SM, Walker M, Rogers AA, Chen WYJ. Moisture balance: optimising the wound-dressing interface. *J Wound Care*, 2003;12:125-128. 10. Bowler P, Jones S, Towers V, Booth R, Parsons D, Walker M. 2010. Dressing conformability and silver-containing wound dressings. *Wounds UK* 6: 14-20. 11. Barnes Y, Amir A, Leshem D, et al. Clinical comparative study of Aquacel and paraffin gauze dressing for split-skin donor site treatment. *Ann Plast Surg*. 2004;53(2):132-136. 12. Kogan L, Moldavsky M, Szvalb S, Govin-Yehudain J. Comparative study of Aquacel and Silvercol treatment in burns. *Ann Burns Fire Disasters*. 2004;17(4):201-207. 13. Brunner U, Eberlein T. Experiences with hydrofibres in the moist treatment of chronic wounds, in particular of diabetic foot. *VASA*. 2000;29(4):253-257. 14. Armstrong SH, Ruckley CV. Use of a fibrous dressing in exuding leg ulcers. *J Wound Care*. 1997;6(7):322-324. 15. Piaggini A, Baccetti F, Rizzo L, Romanelli M, Navalesi R, Bonzi L. Sodium carboxyl-methyl-cellulose dressings in the management of deep ulcerations of diabetic foot. *Diabet Med*. 2001;18(4):320-324. 16. Clarke JN, Deakin AH, Dillon JM, Emmerson S, Kinninmonth AWG. A prospective clinical audit of a new dressing design for lower limb arthroplasty wounds. *J Wound Care*. 2009;18(1):5-11. 17. Cohn SM, Looze PP, Brown M, et al. University of Miami Wound Study Group. Open surgical wounds: how does Aquacel compare with wet-to-dry gauze? *J Wound Care*. 2001;13(1):10-12. 18. Ravenscroft MJ, Harker J, Buch KA. A prospective, randomised, controlled trial comparing wound dressings used in hip and knee surgery Aquacel and Tegaderm versus Occlusi-Adh. *Ann R Coll Surg Engl*. 2006;88(1):16-22. 19. Coutts P, Sibbald RG. 2005. The effect of a silver-containing Hydrofiber® dressing on superficial wound bed and bacterial balance of chronic wounds. *Int J Wound J*; 2: 348-356. 20. Jurejak F, Drape T, Jchrastone A, Ofken T, Wyowie Z, Hollander D. 2007. Randomised clinical trial of Hydrofiber dressing with silver versus povidone-iodine gauze in the management of open surgical and traumatic wounds. *J Wound J*; 4: 66-76. 21. Blome-Eberwein, Johnson RM, Miller SF, Caruso DM, Jordan MH, Milner S, Tiedget EE, Sittig KM, Smith L. 2010. Hydrofiber dressing with silver for the management of split-thickness donor sites: A randomised evaluation of two protocols of care. *Burns*; 36: 665-672. 22. Jude EB, Apelqvist J, Sprault M, Martini J. 2004. Prospective randomised controlled study of Hydrofiber dressing containing Iodine silver or calcium alginate dressings in non-ischaemic diabetic foot ulcers. *Diabetic Medicine*; 21: 280-288. 23. Caruso DM, Foster KN, Hermens MHE, Rick C. 2004. AQUACEL™ Ag in the management of partial-thickness burns: Results of a Clinical Trial. *J Burn Care Rehabil*; 25: 89-97. 24. Kwon Lee S. 2003. Clinical experiences with technologies: case reports on the use of two Hydrofiber® dressings. *Ostomy Wound Management*; 49: (8) Suppl. 6-9. 25. Lohane P, Potokar TS. 2006. AQUACEL™ Ag in paediatric burns – a prospective audit. *Annals of Burns and Fire Disasters*; XIX (3): 144-147. 26. Caruso DM, Foster KM, Blome-Eberwein SA, Twomey JA, Herdon DN, Luterman A, Silverstein P, Antimartino JR, Bauer GJ. 2006. Randomized clinical study of Hydrofiber dressing with silver or silver sulfadiazine in the management of partial-thickness burns. *J Burn Care Res*; 27: 298-306. 27. Huang S-H, Yang P-S, Wu S-H, Chang K-P, Lin T-M, Lin S-D, Lai C-S, Lee S-S. 2010. AQUACEL™ Ag with Vaseline gauze in the management of toxic epidermal necrolysis (TEN). *Burns*; 36: 121-126. 28. Harroing KG, Price P, Robinson B, Thomas S, Helfman D. Cost and dressing evaluation of hydrofiber alginate dressings in the management of community-based patients with chronic leg ulceration. *Wounds*. 2001;13(12):229-236. 29. Dillon JM, Clarke JN, Emmerson S, Kinninmonth AWG. The Jubilee method: a modern dressing design which reduces complications and is cost effective following total hip and knee arthroplasty. Poster presented at: American Academy of Orthopaedic Surgeons Annual Meeting; February 14-17 2007; San Diego, CA. 30. Moore PJ, Foster L. Cost benefits of two dressings in the management of surgical wounds. *Br J Nurs*. 2000;9(17):1128-1132. 31. Paddock HN, Fabia R, Giles S, Hayes J, Lowell W, Adams D, Besser GE. 2007. A silver-impregnated antimicrobial dressing reduces hospital costs for pediatric burn patients. *J Pediatric Surg*; 42: 211-215. 32. Muangman P, Pundee C, Opananon S, Muangman S. 2010. A prospective, randomized trial of silver containing hydrofiber dressing versus 1% silver sulfadiazine for the treatment of partial thickness burns. *Int Wound J*; 7: 271-276. 33. Salas S, Tsai C, Chai P. 2009. Clinical evaluation comparing the efficacy of AQUACEL™ Ag Hydrofiber® dressing versus petrolatum gauze with antibiotic ointment in partial thickness burns in a pediatric burn center. *J Burn Care Res*; 30: 380-385. 34. Koyuncu A, Karadag H, Kurt A, Aydin C, Topcu O. 2010. Silver-impregnated dressings reduce wound closure time in marsupialized pilonidal sinus. *EWMA Journal*; 10: 25-27. 35. Preliminary assessment of the physical properties of AQUACEL™ EXTRA vs AQUACEL™. *Scientific Background Report*. WHR13461 TA214 Rev. 1 2011. Data on file, ConvaTec Inc. 36. Bugedo A, Bowler F, Bishop SM. 2012. Assessment of the in vitro physical properties of AQUACEL™ Ag Extra™ and AQUACEL™ Ag dressings. *Technical Assessment*. WHR13602 TA235. 2012. Data on file, ConvaTec Inc. 37. Jones SA, Bowler PG, Walker M, Parsons D. 2004. Controlling wound bioorder with a novel silver-containing Hydrofiber® dressing. *Wound Rep. Regen*. 12: 288-294. 38. Bowler PG, Jones SA, Walker M, Parsons D. 2004. Microbiological properties of a silver-containing Hydrofiber® dressing against a variety of burn wound pathogens. *J Burn Care Rehabil*. 25: 192-196. 39. Antimicrobial activity of AQUACEL™ Ag against superbugs using a seeded agar. *Scientific Background Report*. WHR 13499 MA 180. 2011 Data on file, ConvaTec. 40. Sustained Activity of an antimicrobial Hydrofiber® dressing against antibiotic-resistant wound pathogens P. G. Bowler MPhil., S. A. Jones BSc., M. Walker PhD., D. Parsons PhD., Wound Healing Research Institute, ConvaTec Global Development Centre, Dareside, UK

®/™ sono marchi di proprietà ConvaTec, AQUACEL, AQUACEL EXTRA, HYDROFIBER e relativi loghi sono marchi commerciali registrati negli US.
©2012 ConvaTec Inc.



Quando le lesioni richiedono di più

EXTRA

RESISTENZA • ASSORBIMENTO • AFFIDABILITA'

AQUACEL™ Extra™

AQUACEL™ Ag Extra™

AQUACEL™ | AQUACEL™ Ag | AQUACEL™ Foam
AQUACEL™ Surgical | AQUACEL™ Ag Surgical | AQUACEL™ Burn | AQUACEL™ Ag Burn



AQUACEL™ Dressings
TRIED. TRUE. TRUSTED.™



AQUACEL™
Extra™


AQUACEL™ Ag
Extra™

Per maggiori informazioni visita il sito www.convatec.it o chiama il numero 800.930.930

Le sfide di ogni giorno

- Assorbire e gestire l'essudato aiutando a prevenire il rischio di macerazione¹
- Rimuovere dalla lesione batteri ed enzimi nocivi che possono ritardare il processo di guarigione¹
- Ridurre il dolore e il disagio per il paziente quando la medicazione è in situ o durante il cambio¹
- Ottimizzare i costi garantendo un trattamento efficace¹


Affidati alla Tecnologia Hydrofiber™



Trattiene*

l'essudato della lesione, intrappola i batteri²⁻⁵, riduce la propagazione laterale prevenendo la macerazione della cute perilesionale.^{6,7}

* Come dimostrato *in vitro*.
Tutte le immagini sono rappresentazioni artistiche



Si micro-conforma*

al letto della lesione⁸ mantenendo un ambiente umido bilanciato,⁹ riducendo al minimo gli spazi vuoti dove i batteri possono proliferare.^{8,10}



Interagisce*

con l'essudato della lesione, formando un gel coesivo ed aiutando a ridurre il dolore associato ai cambi di medicazione.^{11,12}

Il valore dell'evidenza clinica

I prodotti della famiglia **AQUACEL™** hanno un patrimonio clinico dal 1995

- Più di 100 pubblicazioni scientifiche di cui 23 studi randomizzati controllati
- Efficacia nel gestire il processo di guarigione¹³⁻²⁷
- Riduzione dei costi generali di trattamento²³⁻³⁴

I benefici della Tecnologia Hydrofiber™ ancora più evidenti in AQ™ Extra™ e AQ™ Ag™ Extra™^{35,36}

Più Tecnologia Hydrofiber™ per un maggior assorbimento e per una migliore gestione dell'essudato

Fibre rinforzate per un extra resistenza

Disponibile con argento

AQUACEL™ Ag Extra™ con l'aggiunta di ioni argento svolge un'efficace attività battericida.³⁷⁻³⁹

Ritagliabile in entrambi i sensi



AQUACEL™ Extra™

39%

più
assorbente^{*35}
e

9x

più resistente di
AQUACEL™^{*35}

AQUACEL™ Ag Extra™

50%

più
assorbente^{*36}
e

9x

più resistente di
AQUACEL™ Ag^{*36}

L'aggiunta di ioni d'argento garantisce una rapida, ed efficace attività antimicrobica. *

- Il serbatoio di ioni argento contenuto nella medicazione, garantisce un'azione battericida contro un ampio spettro di batteri in modo rapido³⁷ (30 minuti) e duraturo⁴⁰ (fino a 14 giorni)
- Ha un'azione battericida³⁷⁻³⁹ su un'ampio spettro di patogeni inclusi MRSA, VRE e germi resistenti agli antibiotici

* Come dimostrato in vitro